

Geometric Mechanics and the Dynamics of Asteroid Pairs

Wang-Sang Koon, Jerrold E. Marsden and Shane Ross
Control and Dynamical Systems 107-81
Caltech, Pasadena, CA 91125

Martin Lo
Navigation and Mission Design
Jet Propulsion Laboratory
California Institute of Technology, M/S 301-140L
4800 Oak Grove Drive, Pasadena, CA 91109

Dan J. Scheeres
Department of Aerospace Engineering
University of Michigan
Ann Arbor, MI 48109-2140

Conference on *New Methods in Astrodynamics*, January, 2003

Abstract

The purpose of this paper is to describe the general setting for the application of techniques from geometric mechanics and dynamical systems to the problem of asteroid pairs. It also gives some preliminary results on transport calculations and the associated problem of calculating binary asteroid escape rates. The dynamics of an asteroid pair, consisting of two irregularly shaped asteroids interacting through their gravitational potential is an example of a *full body problem* or FBP in which two or more extended bodies interact.

One of the interesting features of the binary asteroid problem is that there is coupling between their translational and rotational degrees of freedom. General FBPs have a wide range of other interesting aspects as well, including the 6-DOF guidance, control, and dynamics of vehicles, the dynamics of interacting or ionizing molecules, the evolution of small body, planetary, or stellar systems, and almost any other problem where distributed bodies interact with each other or with an external field. This paper focuses on the specific case of asteroid pairs using techniques that are generally applicable to many other FBPs. This particular full 2-body problem (F2BP) concerns the dynamical evolution of two rigid bodies mutually interacting via a gravitational field. Motivation comes from planetary science, where these interactions play a key role in the evolution of asteroid rotation states and binary asteroid systems.

The techniques that are applied to this problem fall into two main categories. The first is the use of geometric mechanics to obtain a description of the reduced phase space, which opens the door to a number of powerful techniques such as the energy-momentum method for determining the stability of equilibria and the use of variational integrators for greater accuracy in simulation. Secondly, techniques from computational dynamical systems are used to determine phase space structures important for transport phenomena and dynamical evolution.

Contents

1 Introduction	2
2 The F2BP and Asteroid Binaries	4
3 Phase Space Structures	13
4 A Simplified Model for Binary Asteroids	13
5 Conclusions and Future Directions	19

1 Introduction

Full body problems (FBPs) are concerned with the dynamical interaction between two or more distributed bodies. This fascinating class of problems has many interesting open questions and touches on numerous important issues in science and engineering. Examples include binary asteroids, the evolution and dynamics of the Earth-Moon system, the dynamics and control of a high-performance aircraft, reaction and ionization of molecules, interactions and collisions between galaxies, stability and control of underwater vehicles, rendezvous and docking of space vehicles, fine pointing control of a space-based telescope, etc. Geometric mechanics and dynamical systems theory along with appropriate computational and geometric control techniques, provide a unified approach to the analysis and simulation of these problems.

There are many examples in which geometric mechanics methods have been used for this general class of problems. Two examples are the work concerning the use of reduction and stability methods for rigid bodies with flexible attachments¹ and also work on the use of the energy-momentum method for tethered satellites.²

The Binary Asteroid Problem. In this paper we focus on the dynamics of binary asteroids in the context of F2BPs. This problem concerns the dynamics of two spatially extended bodies that interact via their mutual gravitational fields. An example of a motivating goal is the accurate estimation of ejection, collision, and transport rates, accounting fully for coupling between the rotational and translational states of the bodies. We put this problem into the context of systematic approaches to determine the stability of relative equilibria as well as the computation of phase space structures, such as periodic orbits, quasi-periodic orbits, and the division of phase space into regions of regular and chaotic motion. This context allows one to bring to bear powerful computational transport techniques, such as set oriented methods and lobe dynamics. As we shall review below, there is already quite a bit known about relative equilibria in the binary asteroid problem.³ However, rather little has been done on the energy-momentum method (and its converse) for relative equilibria as well as the problem of merging geometric mechanics and dynamical systems calculations, including phase space structure. Our goal is

to take some first steps in this direction, but much work remains to be done on this problem.

The binary asteroid problem is of considerable astrodynamical interest. For instance, the methods are directly relevant to asteroid rotational evolution, variation of planetary obliquities, and the long-term dynamics of Kuiper and asteroid belt binaries, including an analysis of the Pluto-Charon system.

A Little Biased History. Special cases of the F2BP have been analyzed extensively in the literature, with most applications focusing on the interaction of a small, distended body with a much larger body, for example the evolution of irregularly shaped planetary moons or the dynamics and control of gravity gradient spacecraft. Even for this simplified version of the F2BP it is well-known that chaotic motion exists and can play an essential role in the life of a tidally evolving body.^{4,5} Studies from the astrodynamics community have discovered many fundamental symmetries that exist in these ideal problems,^{6,7,8,9} which have in turn been used to develop novel methods for the control of these systems.^{10,11,12} The other extreme case of the F2BP has also been investigated, namely the motion of a massless particle in the field of a strongly non-spherical gravity field, with main application to the motion of spacecraft and ejecta relative to an asteroid, comet or moon.^{13,14,15,16,17,18,19}

An early paper on the FBP was by Duboshin,²⁰ although his subsequent studies focused on the necessary force laws needed for certain solutions and symmetries to exist. Maciejewski²¹ gives a modern statement of the F2BP is given, along with a preliminary discussion of relative equilibria possible for these systems, but with no investigation of the dynamical evolution and stability of such problems. More recently, sharp conditions for the Hill and Lagrange stability of the F2BP have been found²² and numerical and analytical investigations of rotational and translational coupling in these problems have been studied.^{23,24,25} These most recent works serve as one starting point for the current paper.

Tools for the FBP. The F2BP is a magnificently symmetric problem with the overall symmetry group being $SE(3)$, the group of Euclidean motions of three space, together with any symmetries of the bodies themselves. Thus, it is ideal for using the tools of geometric mechanics for systems with symmetry, variational integrators, and transport and dynamical systems theory. In this paper we shall take some small initial steps in this direction for simplified, but still non-trivial, versions of the F2BP. It is our hope that this effort will help merge work from a number of groups into a single coherent theory. Some of the salient features of these methods are as follows.

1. **Geometric Mechanics and Reductions of the F2BP.** As hinted at and explained already,^{26,27,28} geometric mechanics has had enormous successes in many areas of mechanics. Previous investigations in the FBP, however, have been accomplished in a fragmentary way and have missed using, for example, the powerful energy-momentum method and its converse for relative equilibria, as well as the use of geometric phases. For example, the converse of the energy-momentum method^{29,30} allows one to study the destabilizing effect of

dissipation and body deformations for relative equilibria that are gyroscopically stabilized on the linear level (that is, one has a saddle point in the augmented energy, but the system eigenvalues are on the imaginary axis, so one cannot easily infer stability or instability in the nonlinear system).

2. **Variational Integrators for the F2BP.** A novel computational technique that is ripe for use on the FBP is that of AVIs, or Asynchronous Variational Integrators. These are numerical simulation methods that allow different time steps at different spatial points and yet have all the advantages of the usual symplectic integrators used in dynamical astronomy.^{31,32} For example, in many FBPs there are large differences in time scales for the dynamics of rotational and translational motions. AVIs, which allow for such adaptation, greatly improve computational efficiency as has been demonstrated in related problems,^{33,34} and hence permit new problems to be directly simulated for longer times and to higher accuracy than previously achieved.
3. **Phase Space Structure and Transport Calculations.** A third key tool that can be brought to bear on this problem concern advances in the computation of transport rates using transition state theory borrowed from chemistry and lobe dynamics from fluid dynamics, where recent computational advances have enabled significant capabilities using these approaches.³⁵ This may be combined with set oriented methods involving concepts from graph theory such as the notion of graph partitions, congestion, almost invariant sets and the Peron-Frobenius operator.³⁶ Drawing on recent advances in computing phase space objects such as periodic orbits and invariant tori for astrodynamical problems, it would be interesting to investigate the link between periodic orbits, resonances, and restrictions (pinch points) in the phase space of the F2BP, structures important for the computation of transport, such as ejection and collision rates.

There are a number of fundamental questions in planetary science that can be addressed by the techniques that we outline, including: Comet nucleus rotational and translational evolution due to outgassing; Long-term simulation and evaluations of the Yarkovsky effect on the rotational and translational motion of asteroids; Tidal evolution of asteroid and Kuiper belt binaries including dissipation and external forces; Dynamical evolution of galaxies that undergo a close approach.

2 The F2BP and Asteroid Binaries

While problems involving rotational and translational coupling populate many areas of science and engineering, for tractability in this paper we will draw on examples that arise in the field of planetary science and dynamical astronomy, as there are many FBPs in this field whose complete understanding is still lacking, and whose simulation depends on gross physical approximations to certain aspects of their motion. Examples include the evolution of the obliquities of the terrestrial planets,^{37,5}

the rotational dynamics of Europa,³⁸ the effect of the Yarkovsky force on translational and rotational evolution of asteroids,^{39,40,41,42} and the evolution of comet nucleus rotation and translation due to outgassing,^{43,44} to name a few. In each of these cases interactions between rotation and translation are important, but detailed simulation and evaluation of these couplings over long time spans is beyond current capability. More importantly, the specialized force environments and physical effects unique to each of these systems has made it difficult to formulate a general, unifying approach to the analysis and simulation of these problems.

Recent studies of the dynamics and evolution of binary asteroids, provides a class of problems that can serve as fundamental, or “canonical,” models that are key stepping stones to a unified approach to FBPs. This class of problems are of interest in their own right, as recent analysis indicates that up to 20% of Near-Earth Asteroids are binaries,⁴⁵ along with the current boom of binary discoveries in the Main Asteroid Belt and the Kuiper Belt^{46,47} (see Figure 2.1). The analysis and modeling of binary asteroids (or similarly binary Kuiper Belt objects) does not allow the usual assumptions of small gravity coefficients, near equilibrium conditions, or long-term stability of motion made in classical approaches to FBPs. Rather, to properly understand the dynamics and evolution of these systems requires a generalized approach to the problem that incorporates these difficulties into its nominal problem statement.




Figure 2.1: Dactyl in orbit about Ida, discovered in 1994 during the Galileo mission,⁴⁸ and 1999 KW4 radar images of its companion, discovered 2001.⁴⁹

Recent Advances. The general F2BP has been studied under many different approximations, but its general statement has only been well-posed relatively recently,²¹ followed by advances in understanding its Hill and Lagrange stability²² and observations on the role of these interactions on the evolution of small body rotation states.^{50,23,25} The coupled motion in this problem can lead to profound dynamics,²⁵ illustrated in Figure 2.2 showing the evolving orbit radius (a) and rotation period (b) of a sphere and tri-axial ellipsoid of equal mass interacting through gravity alone. The total energy of this system is slightly negative, meaning that the two bodies can never mutually escape (that is, they are Hill stable). The system is also

stable against impact (Lagrange stable), and so will evolve *ad infinitum* unless some small external perturbation (such as from the sun) boosts its energy sufficiently to allow for escape, which would also leave the ellipsoid with an extremely slow rotation rate. The phase space of this apparently chaotic system can be tightly constrained using integrals of motion, and separated regions of phase space can be identified as a function of resonances between the system’s rigid body rotation and orbital rotation rates. Figure 2.2(c)²⁵ shows an example of the allowable phase space of this system, with additional restrictions on motion in the phase space, identified as “pinch points” in the figure, that arise from these resonances. We conjecture that such pinch points can be understood via the reduction process from geometric mechanics, and that these will be important in understanding transport and ejection rates.




Figure 2.2: Time history of the orbit radius (a) and rotation period (b) for a gravitationally interacting sphere and tri-axial ellipsoid of equal mass. Poincaré map (c) showing constraints on the eccentricity of the evolving orbit, including pinch points that isolate regions of phase space.

Models In Use. The equations of motion for the F2BP model may be explicitly found and are discussed in a number of references.^{21,22} A widely used model can be

derived when one of the bodies is a massive sphere. The rotational dynamics of the sphere decouples from the system, which still has coupling of rotational and translational dynamics between the non-spherical body and the relative orbit between the two. The normalized and symmetry reduced equations are

$$\begin{aligned} \mathbf{r}'' + 2\boldsymbol{\omega} \times \mathbf{r}' + \boldsymbol{\omega}' \times \mathbf{r} + \boldsymbol{\omega} \times (\boldsymbol{\omega} \times \mathbf{r}) &= \frac{\partial \mathcal{U}}{\partial \mathbf{r}} \\ \mathcal{I} \cdot \boldsymbol{\omega}' + \boldsymbol{\omega} \times \mathcal{I} \cdot \boldsymbol{\omega} &= -\mu \mathbf{r} \times \frac{\partial \mathcal{U}}{\partial \mathbf{r}}, \end{aligned} \quad (2.1)$$

where

$$\begin{aligned} \mathcal{K} &= \frac{1}{\mu} \mathbf{A} \cdot \mathcal{I} \cdot \boldsymbol{\omega} + \mathbf{A} \cdot \mathbf{r} \times (\mathbf{r}' + \boldsymbol{\omega} \times \mathbf{r}) \\ \mathcal{E} &= \frac{1}{2} (\mathbf{r}' + \boldsymbol{\omega} \times \mathbf{r}) \cdot (\mathbf{r}' + \boldsymbol{\omega} \times \mathbf{r}) + \frac{1}{2\mu} \boldsymbol{\omega} \cdot \mathcal{I} \cdot \boldsymbol{\omega} - \mathcal{U} \end{aligned} \quad (2.2)$$

and where $\boldsymbol{\omega}$ is the rotational velocity vector in the body-fixed frame, \mathbf{r} is the relative position vector in the body-fixed frame, \mathbf{A} is the attitude tensor of the non-spherical body, \mathcal{I} is the specific inertia tensor of the non-spherical body, \mathcal{U} is the gravitational force potential of the non-spherical body, and \mathcal{K} and \mathcal{E} are the angular momentum and energy integrals, respectively, of these equations.

The free parameters of the system are the mass distribution of the non-spherical body (normalized by its largest dimension) and the mass fraction between the two bodies, $\mu = M_1/(M_1 + M_2)$ (the same parameter found in the restricted 3-body problem), where M_1 is the mass of the “sphere” and M_2 is the mass of the distributed body. The case when $\mu \rightarrow 1$ corresponds to the motion of a massless distributed body about a point-mass, with an application to a satellite in orbit about a planet. It is important to note that the angular momentum and energy integrals still apply to this problem. The case when $\mu \rightarrow 0$ corresponds to the motion of a material point in the gravity field of the distributed body, with the main application to particle dynamics about an asteroid. In this case, we see that the energy and angular momentum integrals are dominated by the rotational dynamics of the distributed body, and that the contribution of the spherical body’s motion decouples from these integrals.

In this paper we shall use equation (2.1), often with some further simplifications. These simplifications are made to allow for an abbreviated discussion of results, and will be relaxed in future papers. The main simplification made is the restricted, uniformly rotating approximation which occurs when $\mu \rightarrow 0$ and the distributed body (which now has all the mass of the system) is in principal axis rotation about its maximum moment of inertia (assumed to lie along the z -axis). When needed, we will refer to this system as the Restricted Full 2-Body Problem (RF2BP), and we note its similarity to the celebrated restricted 3-body problem, including the existence of a Jacobi integral. The RF2BP, while highly idealistic, is important for understanding the general and qualitative properties of motion about distended bodies in uniform rotation⁵¹ and approximately models the motions of some binary asteroids.

Carrying out such detailed calculations over the time scales of interest for the F2BP is challenging and requires methods designed to conserve the symmetries inherent in the problem and to handle the multiple time scales in FBPs. Additionally, the existence and prediction of the restricted regions of phase space far from an equilibrium point is an example of a complex, poorly understood dynamical phenomenon. The techniques discussed below take some first steps to address this deficiency. The F2BP is ideally suited for the application of geometric mechanics with symmetry, variational integrators, and transport and dynamical systems theory. Due to the generality of its statement, the F2BP can be extended to cover other systems of interest in planetary science, dynamical astronomy, astrodynamics, and chemical dynamics, briefly discussed later in the paper.

Geometric Mechanics and the F2BP. We ultimately want to carry out an analysis of the F2BP by first looking at the problem’s symmetry. The dynamics of the F2BP are invariant under application of orientation preserving rotations and translations of 3-dimensional Euclidean space, i.e., under the application of the symmetry group $SE(3)$. Thus, in the terminology of geometric mechanics, we say that the F2BP carries the symmetry of invariance with respect to $SE(3)$. A well-known property of mechanical systems like the F2BP is that whenever there is a symmetry, there are corresponding conserved quantities.

As the symmetry group becomes larger, we expect that one recovers all the special cases studied in the literature. For example, if one body is an irregular body and the other is cigar-like, the symmetry group is $SE(3) \times S^1$. Relative equilibria and their stability can be studied using the powerful *energy-momentum method*.^{52,28,30} Relative equilibria are key ingredients in identifying and characterizing the possible final states that an evolving binary asteroid can reach and they play a similar role as the libration points in the 3-body problem. Moreover, a systematic geometric mechanical approach will enable the use of modern numerical algorithms, such as variational integrators, in the accurate computations of relevant long term statistics and transport.^{53,33,34}

Symmetry Reduction. The use of geometric and dynamical techniques to study Lagrangian or Hamiltonian mechanics has been enormously successful in a wide variety of engineering and astrodynamical problems, such as the use of the energy-momentum method for stability of satellites with internal rotor controls³⁰ and in heteroclinic and resonance structures in the 3-body problem.⁵⁴ Geometric mechanics starts with the usual formulation of Lagrangian mechanics using variational principles, and Hamiltonian mechanics using Poisson and symplectic geometry. Lagrangian reduction by a symmetry group corresponds to finding reduced variational principles, whereas reduction on the Hamiltonian side corresponds to constructing appropriate reduced Poisson and symplectic structures.

The reduction of a system with configuration manifold Q and Lie symmetry group G occurs at two levels, the first of which corresponds to identifying solutions that are related by a symmetry group motion, and this corresponds to deriving equations of motion on the quotient space TQ/G (for Lagrangian reduction), and

on T^*Q/G , (for Hamiltonian Reduction). Symmetry gives an associated Noether theorem, namely conservation of a momentum map, so that the dynamics are constrained to a momentum surface in the absence of external forcing and dissipation. When appropriate, imposing this constraint explicitly in the equations of motion, yields a further reduction in the dimensionality of the reduced equations, which corresponds to (nonabelian) Routh and symplectic reduction.

$$\begin{array}{ccc}
 \begin{array}{c} \text{Lagrangian Reduction} \\ TQ/G \cong T(Q/G) \oplus \tilde{\mathfrak{g}} \end{array} & \xrightarrow{\mathbb{F}l} & \begin{array}{c} \text{Hamiltonian Reduction} \\ T^*Q/G \cong T^*(Q/G) \oplus \tilde{\mathfrak{g}}^* \end{array} \\
 \uparrow & & \uparrow \\
 \begin{array}{c} \text{Routh Reduction} \\ J_L^{-1}(\mu)/G_\mu \cong T(Q/G) \times_{Q/G} Q/G_\mu \\ \text{Variational Principles} \end{array} & \xrightarrow{\mathbb{F}\hat{R}^\mu} & \begin{array}{c} \text{Symplectic Reduction} \\ J^{-1}(\mu)/G_\mu \cong T^*(Q/G) \times_{Q/G} \tilde{\mathcal{O}}_\mu \\ \text{Symplectic/Poisson Geometry} \end{array}
 \end{array}$$

On the Lagrangian side, general theorems give the structure of the quotient space to be

$$TQ/G \cong T(Q/G) \oplus \tilde{\mathfrak{g}},$$

where \mathfrak{g} is the Lie algebra of G and where $\tilde{\mathfrak{g}}$ is an associated bundle over *shape space* Q/G , as shown in⁵⁵. There is a similar structure theorem on the Hamiltonian side. When the momentum constraints are imposed, on the Hamiltonian side one gets well known cotangent bundle reduction theorems²⁸ while on the Lagrangian side one gets a far reaching, and surprisingly recent, generalization of the classical Routh procedure.⁵⁶ The general scheme of the reduction procedure is shown in the above figure. For the F2BP, the choice of Q and G are easily done and are given in the next section.

Reduction for the F2BP. For the F2BP, the configuration space is $Q = SE(3) \times SE(3)$. Denote material points in a reference configuration by X_i , and the points in the current configuration by x_i . Given $((A_1, r_1), (A_2, r_2)) \in SE(3) \times SE(3)$, points in the reference and current configurations are related by $x_i = A_i X_i + r_i$, $i = 1, 2$. Using the body angular velocity notation defined earlier, the Lagrangian has the standard form of kinetic minus potential energy:

$$\begin{aligned}
 L(A_1, r_1, A_2, r_2) &= \frac{1}{2} \int_{\mathcal{B}_1} \|\dot{x}_1\|^2 d\mu_1(X_1) + \frac{1}{2} \int_{\mathcal{B}_2} \|\dot{x}_2\|^2 d\mu_2(X_2) \\
 &\quad + \int_{\mathcal{B}_1} \int_{\mathcal{B}_2} \frac{G d\mu_1(X_1) d\mu_2(X_2)}{\|x_1 - x_2\|} \\
 &= \frac{m_1}{2} \|\dot{r}_1\|^2 + \frac{1}{2} \langle \Omega_1, I_1 \Omega_1 \rangle + \frac{m_2}{2} \|\dot{r}_2\|^2 + \frac{1}{2} \langle \Omega_2, I_2 \Omega_2 \rangle \\
 &\quad + \int_{\mathcal{B}_1} \int_{\mathcal{B}_2} \frac{G d\mu_1(X_1) d\mu_2(X_2)}{\|A_1 X_1 - A_2 X_2 + r_1 - r_2\|}.
 \end{aligned}$$

Reducing by Overall Translations and Rotations. The preceding Lagrangian has symmetry group $SE(3)$ using the diagonal left action on Q :

$$(A, r) \cdot (A_1, r_1, A_2, r_2) = (AA_1, Ar_1 + r, AA_2, Ar_2 + r).$$

The momentum map corresponding to this symmetry is the total linear momentum and the total angular momentum. The projection from the configuration space to the shape space $\pi : Q \rightarrow Q/G$, is obtained by transforming to the body frame of the second rigid body.^{21,57,58} The reduction is carried out in stages, by first reducing by translations, \mathbb{R}^3 , followed by reducing by rotations, $SO(3)$. Results from general reduction by stages^{55,59,60} shows that this is equivalent to directly reducing by $SE(3)$ in a single step. This is achieved by applying the inverse of (A_2, r_2) , which is given by $(A_2^T, -A_2^T r_2) \in SE(3)$,

$$\begin{aligned} (A_2^T, -A_2^T r_2) \cdot (A_1, r_1, A_2, r_2) &= (A_2^T A_1, A_2^T r_1 - A_2^T r_2, A_2^T A_2, A_2^T r_2 - A_2^T r_2) \\ &= (A_2^T A_1, A_2^T (r_1 - r_2), e, 0). \end{aligned}$$

Shape space Q/G is isomorphic to one copy of $SE(3)$, being coordinatized by the *relative attitude* $A_2^T A_1 = A^T$ and *relative position* $A_2^T (r_1 - r_2) = R$. The equations of motion in $T(Q/G)$ (resp. $T^*(Q/G)$) involve A, R , and their velocities (resp. conjugate momenta Γ, P), which correspond to total angular and linear momenta in the body fixed frame of the second rigid body. These are coupled to equations in $\mathfrak{se}(3)^*$, which may be identified with equations for the body angular and linear momenta of the second rigid body, Γ_2, P_2 .

The equations for Γ and Γ_2 are rewritten in terms of Γ_1, Γ_2 , and the linear momentum of the second rigid body is ignored.^{21,57,58} A reconstruction-like equation (that is, recovering the full attitude of both bodies) is added at the very end. One thing missing from the literature is the systematic use of shape space to study *geometric phases* that are important for rotational and translational drifts.^{61,59} The general context of geometric mechanics and the specific setting of the present paper should enable one to fill in this interesting gap. The coordinatization of the reduced space by transforming to the reference frame of the second rigid body corresponds to a particular choice of connection on the principal bundle $Q \rightarrow Q/G$. Different choices of connection will affect how we identify TQ/G with $T(Q/G) \oplus \mathfrak{g}$.

The reduced Lagrangian is obtained by rewriting the Lagrangian in terms of the reduced variables:

$$A = A_2^T A_1, \quad R = A_2^T (r_1 - r_2), \quad \hat{\Omega} = A_2^T \dot{A}_1, \quad V = A_2^T (\dot{r}_1 - \dot{r}_2),$$

which are coordinates on $T(Q/G)$, as well as $\hat{\Omega}_2 = A_2^T \dot{A}_2$, $V_2 = A_2^T \dot{r}_2$, which are coordinates on $\mathfrak{se}(3)$. As with the Euler-Poincaré theory,²⁷ Hamilton's variational principle on $T(SE(3) \times SE(3))$ is equivalent to the *reduced variational principle*,

$$\delta \int_a^b l(A, R, \hat{\Omega}, V, \hat{\Omega}_2, V_2) dt = 0,$$

on \mathbb{R}^{18} where the variations are of the form,

$$\begin{aligned} \delta A &= -\hat{\Sigma}_2 A + \hat{\Sigma}, & \delta R &= -\hat{\Sigma}_2 R + S, & \widehat{\delta\Omega} &= \dot{\hat{\Sigma}} - \hat{\Sigma}_2 \hat{\Omega} + \hat{\Omega}_2 \hat{\Sigma}, \\ \delta V &= \dot{S} - \hat{\Sigma}_2 V + \hat{\Omega}_2 S, & \delta\Omega_2 &= \dot{\hat{\Sigma}}_2 + \Omega_2 \times \Sigma_2, & \delta V_2 &= \dot{S} - \hat{\Sigma}_2 V_2 + \hat{\Omega}_2 S_2, \end{aligned}$$

and S, S_2, Σ, Σ_2 are variations that vanish at the boundary. The symplectic form can be obtained from the variational principle by considering variations that do not vanish at the boundary. The boundary terms that arise from integration by parts can be interpreted as a Lagrange one-form, and taking the exterior derivative of these yield the symplectic two-form.

Already the above approach is more systematic and complete than that given in the literature. While simple, the variational structure just given is, in fact, new. In future studies, we will complete this task and deal with all of the systematic cases for the F2BP and to tie in this theory with the stability theory of relative equilibria using the powerful energy-momentum method.

The Energy-Momentum Method. This is a powerful generalization of the classical energy methods for stability.⁵² A study of the stability of rigid body pairs using the energy-momentum method is directly relevant to the evolution of binary asteroids and the dynamical environments they encounter through their lifetime. The current literature, however, uses only the Arnold or energy-Casimir methods⁵⁷ which generally give weaker results. It would be very interesting to make use of the full block diagonalization power of the energy-momentum, which has proven useful in related problems.^{7,2} In addition, there is a converse to the method which allows one to deduce the development of instabilities due to forcing and dissipation,^{29,30} which can have important long term effects on the dynamics.⁶²

Symmetries of the Rigid Bodies. If, in addition, the individual rigid bodies exhibit configurational symmetries of their own, there are additional symmetries acting on the right. Since these commute with the symmetry of $SO(3)$ acting on the left, one can again make use of reduction by stages.

Example: Spherical Symmetry. If the first body is spherical, and the second body has no symmetry, the additional symmetry group is $SO(3) \times \{e\}$ acting on the right. On the Hamiltonian side, the reduced space is then $T^*\mathbb{R}^3 \times \mathfrak{se}(3)^* \times \mathfrak{so}(3)^*$. The $T^*\mathbb{R}^3$ terms correspond to position and linear momentum of the center of mass of the first rigid body, in the body frame of the second rigid body. The $\mathfrak{se}(3)^*$ term involves the spatial linear and angular momentum of the second rigid body, and $\mathfrak{so}(3)^*$ involves the body angular momentum of the first rigid body. There is a similar picture on the Lagrangian side.

Example: Cylindrical Symmetry. If the first body is cylindrical, and the second body has no symmetry, the additional symmetry group is $S^1 \times e$ acting on the right. The reduced space is then $T^*(S^2 \times \mathbb{R}^3) \times \mathfrak{se}(3)^* \times \mathbb{R}$. The $T^*(S^2 \times \mathbb{R}^3)$ terms

will correspond to the orientation of the symmetry axis and the position of the first rigid body, in the body frame of the second rigid body, as well as the conjugate momenta. The $\mathfrak{se}(3)^*$ term involves the spatial linear and angular momentum of the second rigid body, and the \mathbb{R} term will involve the component of the body angular momentum of the first rigid body in the direction of its symmetry axis.

The preceding reduction procedures give a global description of the reduced spaces and dynamics. This is important, since ad hoc choices for coordinate systems which are only locally valid can introduce significant computational overheads if it is constantly necessary to switch between local charts in the course of a long time simulation. The variational, Poisson, and symplectic structures on these reduced spaces can be constructed systematically from the general theory of reduction, and, as we have indicated, this is particularly useful in classifying a hierarchy of simplified models.

Relative equilibria and relative periodic orbits correspond to equilibria and periodic orbits in the shape space Q/G . The bifurcation of a relative equilibria into a relative periodic orbit through a Hamiltonian Hopf bifurcation in the shape space is also particularly interesting for the study of real world objects, which may be subject to parametric uncertainty, and the effects of such parameter dependence need to be studied in our models. More generally, the study of these phase space objects, their stability, and their bifurcation patterns can be used to characterize motion in different regions of phase space.⁶³

The phenomena of pinch points in the dynamics could possibly be related to the presence of resonances corresponding to discrete symmetries, or possibly to degenerate values of the momentum map corresponding to the non-free action of a continuous symmetry group. It is well known that pinch points of this sort are related to reduction in singular cases.⁶⁴ An example of a problem with such a non-free action is the double spherical pendulum, and the bifurcation from the degenerate downward configuration into relative equilibria and relative periodic orbits has been analyzed using the machinery of bifurcation theory and singular reduction theory.⁶⁵ A better understanding of the pinch points observed in dynamical systems will be directly applicable for identifying the regions of phase space which contribute to statistical transport phenomenon.

Geometric Phases. The issue of geometric phases is not only of interest in the dynamics of an asteroid pair, but is also relevant to the problem of landing a spacecraft on a spatially extended and irregular asteroid (as in the recent NEAR mission to the asteroid EROS). An understanding of geometric phases allows one to relate the motion of internal rotors, and their effect on the relative orientation of the spacecraft with respect to the asteroid. This provides a precise method of controlling the relative orientation, which is in contrast to the use of micro-thrusters, which introduce torque which have to be compensated for once the desired orientation is achieved. This same approach can be applied to the study of the tidally locked Pluto-Charon system and may help address some aspects of the current controversy surrounding the value of Charon's orbital eccentricity and how it arose.⁶⁶

Geometric phases can also be applied to spacecraft control in close-proximity

to an asteroid.⁶⁷ In this context of control, the method of Controlled Lagrangian and Hamiltonian systems may be a useful tool.⁶⁸ This is generalization of potential and kinetic shaping techniques which has been applied in the context of spacecraft dynamics to modify the system’s Poisson structure using feedback control laws. A corresponding theory for the reduction of systems with symmetry also exists, using the method of Lagrangian and Hamiltonian reduction.

3 Phase Space Structures

Global Theory of Chaotic Transport. After systematically categorizing the systems of interest using geometric mechanics, the next step is to use efficient computations guided by dynamical systems theory to explore statistical questions. These questions, including the probability of binary pair formation and subsequent escape or impact, and ejecta redistribution around a rotating small body can be cast as *phase space transport* problems. The systems of interest, being Hamiltonian in nature, likely have a global mixed phase space structure of stable and chaotic zones, i.e., islands of KAM tori and a “chaotic sea” between them. A semi-analytical *global theory of chaotic transport* is emerging which combines the theory and numerics of lobe dynamics and tube dynamics.^{69,54,70} The lobe dynamics techniques come partly from fluid dynamics—e.g., those developed in the last two years using the MANGEN software—providing unprecedented long-term precision calculations.^{71,72}

Efficient Computation of Statistical Quantities. This global theory allows one to tackle transport problems by focusing on the objects important for transport. Dynamical systems theory has been used previously to give a qualitative description of the topological features of the phase space in certain reduced F2BP models, such as periodic orbits, resonance regions, and chaotic zones. One can make this analysis more quantitative by computing statistical measures such as residence times within regions of phase space, transport rates between various regions, and the overall level of mixing between regions.

Since the F2BP is such a fundamental model in studying small body orbital and rotational dynamics, the present context should shed light on several interesting and important problems. Specifically, it may help address the problem of slowly rotating asteroids,⁵⁰ it will be able to quantify the rate at which asteroid binaries are disrupted due to their mutual interactions following their creation,²⁵ and the important issue of chaotic planetary obliquities and their probability of transition to different states.³⁷

4 A Simplified Model for Binary Asteroids

The Restricted Full 2-Body Problem (RF2BP). The simplest model that exhibits the basic ejection and collision dynamics we are interested in studying is given by the RF2BP, which is a simplification of Equation (2.1) in the limit $\mu \rightarrow 0$. The equations of motion for the massless particle in a rotating Cartesian coordinate

frame and appropriately normalized are

$$\begin{aligned}\ddot{x} - 2\dot{y} &= -\frac{\partial U}{\partial x} \\ \ddot{y} + 2\dot{x} &= -\frac{\partial U}{\partial y},\end{aligned}\tag{4.1}$$

where

$$\begin{aligned}U(x, y) &= -\frac{1}{\sqrt{x^2 + y^2}} - \frac{1}{2}(x^2 + y^2) + U_{22} \\ U_{22} &= -\frac{3C_{22}(x^2 - y^2)}{(x^2 + y^2)^{5/2}}\end{aligned}$$

The system (4.1) has one free parameter, the gravity field coefficient C_{22} , commonly termed the “ellipticity”, that varies between 0 and 0.05 for physical systems. The system (4.1) may be readily derived from Lagrangian mechanics using the method of “moving systems”²⁷ and therefore has an energy integral (a Jacobi integral), namely

$$E = \frac{1}{2}(\dot{x}^2 + \dot{y}^2) + U\tag{4.2}$$

This system has 4 equilibrium points symmetrically placed along the x and y axes, each at a radius of $R \sim 1 + \dots$, with the higher order terms arising from the U_{22} potential. In general, we do not consider motion at radii much less than $R \sim 1$, since these trajectories will usually impact on the central body. Also, note that these equations have very bad behavior as $R \rightarrow 0$, which is another reason to avoid radii much inside the equilibrium points.

The region of space where $R > 1$ is very interesting, however. First, there are continuous families of near-circular periodic orbits that exist for any given radius. Occasionally, these periodic orbits are resonant with the rotation rate of the system, and at these points we can find additional families of “elliptical” periodic orbits that branch off.

Realms and Regions in the RF2BP. In the RF2BP, the value of the energy E indicates the type of global dynamics possible. For example, for energies above a threshold value, (i.e., $E > E_S$, where E_S corresponds to the energy of symmetric saddle points along the x -axis), movement between the *realm* near the asteroid (*interior realm*) and away from the asteroid (*exterior realm*) is possible, as in Figure 4.1(a). This motion between realms is mediated by phase space *tubes*, to be described shortly. For energies $E \leq E_S$, no such movement is possible. Within each realm, the phase space on appropriately chosen Poincaré sections is organized further into different *regions*, connected via *lobes*.

Tube Dynamics: Transport between Realms. In Figure 4.1(a) the interior and exterior realms are linked by tubes in phase space, bounded by the stable and unstable invariant manifolds associated to periodic orbits around the symmetric




Figure 4.1: **Planar restricted F2BP with uniform rotation: realms and allowable motion.** (a) For energies, E , above a threshold, E_S , movement is possible between the realm near an asteroid and away from the asteroid. In this rotating frame, the x -axis coincides with the elliptical asteroid’s long axis. The origin is shown as a large black dot at the center of the asteroid. The value of the “ellipticity”, C_{22} , is 0.05. Varying this parameter changes the shape of the asteroid and the subsequent potential through the U_{22} term in Equation (4.1). (b) Poincaré sections in different realms—in this case in the *interior* and *exterior realms*, U_1 and U_2 , respectively—are linked by *phase space tubes* which live in surfaces of constant energy (Equation (4.2)). Under the Poincaré map f_1 on U_1 , a trajectory reaches an *exit*, the last Poincaré cut of a tube before it enters another realm. The map f_{12} takes points in the exit of U_1 to the *entrance* of U_2 . The trajectory then evolves under the action of the Poincaré map f_2 on U_2 .

saddle points. The role these tubes play in global transport between realms is referred to as *tube dynamics*.⁵⁴ On each three-dimensional energy surface these two-dimensional tubes partition the surface, acting as separatrices for the flow through the bottleneck regions around the saddle points: particles inside the tubes will move between realms, and those outside the tubes will not. For example, an ejecta particle liberated from the surface of the asteroid into the exterior realm with an energy just above the threshold can impact the asteroid *only* by passing through one or the other of the pair of exterior branch stable manifold tubes associated to the two periodic orbits around the saddle points (the right periodic orbit is shown in Figure 4.1(a)).

The projection of these tubes onto the configuration space appear as strips and trajectories on the tubes wind around them. A few trajectories on a couple of tubes are shown in Figure 4.1(a). On the Poincaré section, the last Poincaré cut of the stable manifold tube is called the *exit*, because points in there will exit the interior realm and go to the exterior realm. The time reverse situation holds for an *entrance*, and when a particle goes through the exit in one realm, it must enter the entrance of another.

These transition dynamics are of direct interest to the evolution of a dissipating binary system, as it may experience several junctures when it can transition between a state close to its final equilibrium and a state where more dynamic evolution is possible. Similarly, the long-term variation of planetary obliquities may be modeled using this approach.

Lobe Dynamics: Transport between Regions. Tubes are only one part of the global transport picture. The study of transport *between regions* within a particular *realm* can be reduced to the study of an associated Poincaré section in that realm. Lobe dynamics theory states that the two-dimensional phase space M of the Poincaré map f can be divided into disjoint *regions* of interest, $R_i, i = 1, \dots, N_R$, such that $M = \cup R_i$ (Figure 4.2(a)).⁷³ The boundaries between regions are pieces of stable and unstable manifolds of hyperbolic fixed points, $p_i, i = 1, \dots, N$. Moreover, transport between regions of phase space can be completely described by the dynamical evolution of *turnstile lobes* enclosed by segments of the stable and unstable manifolds (Figure 4.2(b)).




Figure 4.2: **Transport between regions in phase space.** (a) A point q_k is called a *primary intersection point* (pip) if $S[p_i, q_k]$ intersects $U[p_j, q_k]$ only at the point q_k , where $U[p_j, q_k]$ and $S[p_i, q_k]$ are segments of the unstable and stable manifolds, $W^u(p_j)$ and $W^s(p_i)$ respectively, of unstable fixed points of the Poincaré map f . The union of segments of the unstable and stable manifolds form partial barriers, or *boundaries* $U[p_j, q_k] \cup S[p_i, q_k]$ between *regions* of interest $R_i, i = 1, \dots, N_R$, in $M = \cup R_i$. The region on one side of the boundary B_{12} is labeled R_1 and the other side labeled R_2 . (b) Let $q_0, q_1 \in W^s(p_i) \cap W^u(p_j)$ be two adjacent pips, i.e., there are no other pips on $U[q_0, q_1]$ and $S[q_0, q_1]$, the segments of $W^u(p_j)$ and $W^s(p_i)$ connecting q_0 and q_1 . The region interior to $U[q_0, q_1] \cup S[q_0, q_1]$ is a *lobe*. Then $S[f^{-1}(q_0), q_0] \cup U[f^{-1}(q_0), q_0]$ forms the boundary of two lobes; one in R_1 , labeled $L_{1,2}(1)$, and the other in R_2 , labeled $L_{2,1}(1)$. Under one iteration of f , the only points that can move from R_1 into R_2 by crossing B_{12} are those in $L_{1,2}(1)$ and the only points that can move from R_2 into R_1 by crossing B_{12} are those in $L_{2,1}(1)$. The union of the two lobes, $L_{1,2}(1) \cup L_{2,1}(1)$, is called a *turnstile*.

To keep track of points as they move between regions, suppose that, at $t = 0$, region R_i is uniformly covered with points of *species* S_i . The transport problem

becomes one of describing the distribution of species S_i throughout the regions R_j for any time $t = n > 0$. Two quantities of interest—the flux of species S_i into region R_j on the n -th iterate, $\alpha_{i,j}(n)$, and the total amount of species S_i in region R_j just after the n -th iterate, $T_{i,j}(n)$ —can be expressed compactly in terms of intersection areas of images or pre-images of turnstile lobes. In our application, the Poincaré map of a reduced F2BP model possesses resonance bands consisting of alternating unstable and stable periodic points. For instance, we can study the transport from stable configurations to escape configurations or the transport into and out of spin-orbit resonances, and many other transport questions that have implications for the current state of bodies in the solar system.

Computational Tools for Lobe Dynamics. While lobe dynamics has always been recognized as an *exact* transport theory and can theoretically give short and long term transport rates, computational issues have limited its applications (^{74,75}). The manifolds computed in such problems are typically rather convoluted, as implied in Figure 4.2. Furthermore, the length of these complicated curves grows quickly with the size of the time window of interest. The number of points needed to describe long segments of manifolds can be prohibitively large if brute force approach computational methods are used.

Recent efforts made to incorporate lobe dynamics into geophysical and chemical transport calculations have brought new techniques to compute invariant manifolds.^{76,70,71,77,72} We have been able to compute long segments of stable and unstable manifolds with high accuracy by *conditioning the manifolds adaptively*, e.g., by inserting more points along the manifold where the curvature is high. As a result, the length and shape of the manifold is not an obstacle, and many more iterates of lobes than hitherto possible can be generated accurately. Initial tests show that this method of computing lobes using MANGEN is very relevant to the problems we investigate.

Phase Space Structure in the Exterior Region. A Poincaré surface-of-section (s-o-s) of the RF2BP at an energy just above the threshold where the bottleneck appears (i.e., $E > E_S$, the case in Figure 4.1(a)) illustrates the relevance of tube and lobe dynamics in this system. In Figs. 4.3(a,b), the s-o-s was taken in the exterior region along the positive x -axis. At the energy chosen, there is a bottleneck around the equilibrium points along the x -axis. Particles beginning in the exterior region will reach the interior region and subsequently collide with the asteroid if they lie within the phase space tubes associated with the unstable periodic orbits about either the left or right saddle points. Thus we refer to the tube slices on this s-o-s as *tube slices of collision*. Furthermore, particles will be ejected from the system if they lie within lobes enclosed by the stable and unstable manifolds of a hyperbolic fixed point at $(+\infty, 0)$, referred to as *lobes of ejection*.

Physical insight is gained into the conditions for ejection by transforming to *Delaunay variables*. In Figure 4.3(b), the semimajor axis is shown versus the argument of periapse with respect to the rotating asteroid (the body-fixed frame). The alternate fates of collision and ejection are seen to be intimately intermingled in the




Figure 4.3: Mixed phase space structure for the restricted model: the dynamics of particle ejection from, and collision with, a distended asteroid. A Poincaré surface-of-section taken in the exterior region ($R > 1$). (a) Particles not ejected after 15 iterates are shown. The coordinates are the radial velocity of the particle versus its radial distance from the asteroid’s center. The finger-like structures visible here, the *lobes of ejection*, have been seen in some chemistry problems.^{78,70} (b) Semi-major axis vs. Argument of Periapse with respect to the rotating asteroid (the body-fixed frame). (c) An interesting trajectory corresponding to the labeled point on the Poincaré section in (a) is shown. The trajectory escapes the asteroid only to fall back upon it after one large elliptical orbit. The units are in terms of the semiaxis length of the asteroid, which is shown schematically at one instant of time. The asteroid is rotating counterclockwise around its center (shown as a large dot at the origin). The long axis of the asteroid sweeps out the circular region bounded by the dashed line.

phase space. Note that the number of particles remaining in the fourth quadrant ($270^\circ - 360^\circ$) is smaller than that in the other three quadrants, in agreement with

earlier observations.¹⁶ Figure 4.3(c) shows a particular trajectory which escapes from the asteroid only to fall back upon it, a peculiar phenomenon encountered in tube dynamics.⁵⁴

In the RF2BP, we are considering the motion of a massless particle, i.e., the limit $\mu \rightarrow 0$. The next step in our study is to assume that the particle has mass, i.e., $\mu > 0$ in Eq. 2.1. Due to the gravitational attraction of the particle, the rotation of the asteroid is now non-uniform and must be tracked. While this adds another dimension to the phase space, it also adds a new integral, the angular momentum \mathcal{K} in Eq. (2). Thus we can still use the two-dimensional Poincaré map analysis, as in Figure 4.3. We are currently in the process of characterizing the global phase space structure which gives rise to the complex behavior seen in Figure 2.2 and similar problems.

5 Conclusions and Future Directions

We have outlined the general setting for the application of techniques from geometric mechanics and dynamical systems transport calculations to *full body problems* (FBPs). General FBPs have a wide range of other interesting aspects as well, including the 6-DOF guidance, control, and dynamics of vehicles, the dynamics of interacting or ionizing molecules, the evolution of small body, planetary, or stellar systems, and almost any other problem where distributed bodies interact with each other or with an external field.

This paper focused on a motivating example of a full 2-body problem from dynamical astronomy—the problem of asteroid pairs and to the calculation of binary asteroid escape rates. We have given some preliminary results for a simplified model of binary asteroid escape rates, describing how lobe and tube dynamics lead to a rich phase space even for this simplified system.

Some future directions for this area of research are as follows.

Variational Integrators for FBPs. As is noted in the literature, the Poisson structures that are obtained in the reduced models are non-canonical, and as such applying standard symplectic algorithms will not provide the long-time stability we have come to associate with such numerical methods, since they preserve the canonical symplectic form, which is not consistent with the reduced dynamics. Instead, we shall to make use of reduction theory for discrete systems, the abelian Routh case of which is worked out and applied to the J_2 problem (satellites in orbit about an oblate planet).⁷⁹ We will study transport phenomena in the F2BP using numerical schemes that preserve the geometric structures inherent in the system. Toward this end, we will redo the above computations for other F2BP cases and carry out long time accurate time integrations using variational integrators that capture transport rates and the detailed structure of chaotic sets.

Variational integrators provide a systematic and powerful extension of symplectic integrators that have a proven track record in celestial mechanics for long-term integrations.^{31,32,80} The variational setting allows for extensions to partial differen-

tial equations, for asynchronous methods as well as the use of symmetry methods that are central to this approach.

The idea of the variational integration method is to discretize Hamilton’s principle directly rather than discretize the associated Euler–Lagrange equations. The discrete form of Hamilton’s principle then determines the numerical algorithm for that system. The accuracy to which the approximation of Hamilton’s principle is done is reflected in the accuracy of the algorithm itself; however, much more is true. When integrators are designed this way, they have remarkable respect for the geometric mechanics aspects of the problem, including excellent energy behavior, conservation of the symplectic structure and exact conservation of a discrete version of the Noether quantities associated with symmetries. It is perhaps surprising that these methods work well even for dissipative and forced systems. The idea in this case is to discretize the Lagrange–d’Alembert principle rather than Hamilton’s principle.

Many recent references develop and document the success of this methodology.^{81,33,53} Figure 5.1 shows a computation for a particle moving in the plane under a radially symmetric polynomial potential, without and with a small amount of friction. The exact preservation of the conserved quantities, in this case energy, is a natural consequence of the discrete variational principle. In addition, Figure 5.1 illustrates that these methods can handle dissipative systems and get the energy decay rate accurate as measured against a benchmark calculation.⁸² In addition, many statistical quantities, such as temperature and the structure of chaotic invariant sets, are accurately captured by variational integrators.^{33,83}




Figure 5.1: Conservative system (left) and dissipative system (right).

A key feature is the development of AVI’s or *asynchronous variational integrators*³³ which allow one to take different time steps at different spatial points and yet retain all the beautiful structure preserving properties of variational integrators. This is important in FBPs since the different bodies may have vastly different spatial and temporal scales, and so one must adapt the time steps according to these different scales. The AVI approach is a natural way to study multi-scale dynamics without sacrificing structure preservation. Interestingly, this is closely related to

methods used in molecular dynamics (work of Barash and Schlick⁸⁴).

Reduction for Discrete Mechanics. There has been significant work done on reduction theory for discrete variational mechanics. An example is that of the dynamics of a satellite in the presence of the bulge of the Earth, in which many interesting links with geometric phases and computation in the reduced and unreduced spaces are noted.⁷⁹ However, for the F2BP, this basic theory needs to be generalized, a key point being the extension to the case of nonabelian symmetry groups and the generalization beyond the Euler-Poincaré context.⁸⁵ We will undertake such a study and to apply it to the case of the F2BP. There is a cautionary message in the work that numerical experience shows—it is important to fully understand the reduction of discrete variational mechanics, since applying standard numerical algorithms to the reduced equations obtained from continuous reduction theory may not yield the desired results and long term stability may not be respected. A related goal is to combine variational integrators and adaptive manifold conditioning to give a state of the art package to compute manifolds, lobes and turnstiles in a variety of systems, including the F2BP. The use of variational integrators is expected to increase the accuracy of computation of long term statistical quantities, such as transport rates. We also may combine these techniques with those of the Delnitz group using box elimination methods.^{86,36}

The Full N -Body Problem. A natural extension of the F2BP is the problem of multiple distributed bodies interacting with each other, the Full N -Body Problem. For example, it would be interesting to put some earlier work²⁰ into a modern context using the same reduction and simulation theory outlined above. Recent success in applying classical N -Body Problem results to the F2BP^{22,25} indicates that progress can indeed be made in developing this approach.

Incorporation of Deformation and Dissipation Effects. The incorporation of deformation and dissipation effects into one or both of the bodies provides an important path towards understanding full body dynamics in a deeper way. Even if the simplest deformation and dissipation models are incorporated, based on small deformation and linear dissipation assumptions, these effects can have major influence on full body attitude and orbital dynamics. This phenomenon is well known in planetary dynamics⁸⁷ and has also been characterized in the field of space structure dynamics and control.

Models that incorporate attitude, translation, and finite deformation degrees of freedom for a distributed body in a central body gravitational field, have been developed.⁸⁸ Even in this relatively simplified version of a FBP, the presence of deformation and dissipation can provide mechanisms for energy transfer between translation and attitude. They also play important roles in controlled FBPs, where physical or control effects directly influence the deformation and dissipation, and indirectly influence attitude and/or translation.

It is also of interest to extend recent work on the stability and control of satellites with flexible appendages to describing the natural dynamics of deformable bodies

in orbit about a rigid body. As examples, the stability and Hamiltonian structure of a rigid body with attached flexible rod was analyzed.¹ More recently,⁸⁹ induced instabilities in a satellite with momentum wheels with an attached string has been considered. This “radiation induced instability” is related to the dissipation induced instabilities studied earlier.^{29,30} Such instabilities can occur when there is a saddle point in the energy-momentum function but all eigenvalues of the linearized system lie on the imaginary axis, situations which occur in the relative equilibria of many FBPs. These results, when coupled with a recent rediscovery of an elastic solution for tri-axial ellipsoids,⁹⁰ could yield significant enhancements in our current understanding of tidal disruptions and dissipation in small body systems.^{91,92}

Astronomy and Planetary Science. Generalized models of the F2BP have many applications in the field of astronomy and planetary science. In the following we mention a few specific targets for our research, chosen as good candidates for the application of our systematic approach to full body problems.

An important outcome of this phase of analysis could involve the long-term simulation of the effects of the Yarkovsky effect on the translational and rotational motion of asteroids. This effect has recently been recognized to be an important component in describing the evolution and current state of the solar system. While much work has been done on the modeling and prediction of its effects,^{39,40,41,42} long-term simulations of its effects using realistic models is currently not possible. The heart of the difficulty is the delicate relation between an asteroid’s rotation state and the net result of the Yarkovsky force (due to thermal inertia in the body). The suite of simulation tools we are developing will be able to directly address this question, and would unquestionably advance our ability to model and understand this important effect.

A related open question in planetary science is an explanation for the large number of slowly rotating asteroids.^{93,50} Current distribution statistics cannot be accounted for by using the traditional theory of asteroid rotation state evolution. While Yarkovsky forces have been considered as a possible explanation,⁴² a strong case has not been built as of yet. Mutual disruption of binaries under their own gravitational interaction is another potential mechanism for the creation of slowly rotating and tumbling asteroids. To properly address whether this can account for the noted excess of slow rotators a deeper study of ejection probability in binary asteroids must be made, starting from earlier work,^{23,25,50} coupled with an analysis of the likely formation energies of these binaries, drawing on recent work in this area.^{94,95,96,92,97,98}

Related to this study is a complete analysis of the formation and evolution of a binary asteroid or Kuiper belt object, including the effect of energy dissipation and external force perturbations. The simulation tools being developed for the F2BP will be ideal for the simulation of these systems over long time spans, allowing for the development and testing of various hypotheses and constraints on the life of these objects. Based on current theoretical models and results^{99,23,25} a binary will be subject to mutual disruption or impact during its evolution, and only a fraction of binary bodies should survive into a long-term stable state. The resolution

and understanding of this question is completely open, but would be naturally addressed using a FBP approach. As part of this analysis, we will also examine the Pluto-Charon system, currently subject to some controversy based on the apparent eccentricity of their mutual orbit.⁶⁶

Recent re-analysis of the long-term evolution of comet nucleus rotation states accounting for realistic models of outgassing torques on the bodies has shown a remarkably rich set of steady-state states for their rotation states.⁴³ Direct simulation of these effects over long-periods is currently not possible, but may be made attainable from the outcomes of our work. Furthermore, the non-trivial dependence on rotation state on a comet's heliocentric orbit invites a more complete analysis of the coupling of nucleus rotation and its orbital variation due to outgassing, which is the classical problem of comet nucleus orbit mechanics.⁴⁴

An additional, novel idea is to apply the F2BP, with dissipation and deformation effects, to describing the interaction of two or more distributed galaxies. As a first step, this work would link the classical theory of ellipsoidal figures of equilibrium¹⁰⁰ with the effect of a non-collision interaction between two galaxies. Following such an encounter, each of the distributed bodies would feel a near-impulsive change in their energy and angular momentum, due to exchanges between the distributed bodies.²⁴ Treating these stellar collections as FBPs could lead to fruitful new insights and approaches to describe their evolution and dynamics.

Vehicle Dynamics and Control. Control problems arise in FBPs where one of the bodies is an artificial body such as a controlled spacecraft or satellite. There is a vast literature on attitude control and orbit control of spacecraft, and the success of most space missions has depended on this capability. With only few exceptions, this literature treats attitude control problems and orbit control problems as independent problems. This is reflected in the organization chart of most space agencies and space industries; attitude control groups and orbit control groups operate independently.

The thesis of this area of research, based on the fundamental definition of FBPs, is that attitude and orbit dynamics and control problems should be treated in a unified way. A consequence of this thesis is that control problems for FBPs should be formulated, analyzed, and implemented using prior knowledge of the full body dynamics. Geometric properties such as symmetries and reduction are important not only for analysis of full body dynamics, but are crucial to full body control as well. Most full body control problems are likely to be nonlinear and to require special methods for control design and analysis that are tailored to the full body dynamics.

Nonlinear control methods should provide excellent frameworks for studying full body control problems. These methods include controlled Lagrangian,¹⁰¹ geometric phases,^{61,10} and differential geometric approaches.^{102,10} These approaches have been developed and successfully applied to the control of rigid bodies and multi-body systems, taking into account motion constraints, symmetries, control-actuation assumptions, deformation degrees of freedom, and dissipation. They have been used to construct controllers for specific rigid-body and multi-body spacecraft attitude control problems using thrusters, reaction wheels, proof mass actuators, appendages,

and tethers. By analyzing and describing these systems from a more general vantage, it is often possible to gain additional insight into the known symmetries of the simpler problem, and potentially to discover new symmetries not appreciated before. A similar philosophy can be applied to a re-analysis of vehicle motion in fluid fields, where the existence of the fluid can be idealized as the limiting model of a second body interaction on the vehicle. As these vehicles have strong coupling between rotational and translational motion, and are considered to be extremely difficult analytical problems, any advance in understanding such vehicles would be important.

Chemical Dynamics. The mathematical description of transport phenomena applies to a wide range of physical systems across many scales.^{103,73,104} The recent and surprisingly effective application of methods combining dynamical systems ideas with those from chemistry to the transport of Mars ejecta by several of the co-authors in collaboration with chemists, underlines this point.³⁵ Thus, techniques developed to study transport in the solar system are truly *fundamental and broad-based*. The methods may be applied to diverse areas of study, including fluid mixing^{105,106,107,108,109,76,71} and N -body problems in physical chemistry.¹¹⁰ A basic framework for this theory has already allowed the several of the co-authors to develop a new low-fuel mission concept to explore Jupiter's moons.¹¹¹ Any improvement made in the software in the course of the work outlined above can thus be applied to areas beyond dynamical astronomy.

Acknowledgements. This research was partially supported by a Max Planck Research Award and NSF ITR grant ACI-0204932. The work described in this paper was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

Author Email Addresses : koon@cds.caltech.edu, marsden@cds.caltech.edu, shane@cds.caltech.edu, Martin.Lo@jpl.nasa.gov, scheeres@umich.edu

References

1. Krishnaprasad, P. S. and Marsden, J. E. [1987] Hamiltonian structure and stability for rigid bodies with flexible attachments. *Arch. Rat. Mech. Anal.* **98**:137–158.
2. Krupa, M., Schagerl, M., Steindl, A., et al. [2001] Relative equilibria of tethered satellite systems and their stability for very stiff tethers. *Dyn. Syst.* **16**:253–278.
3. Scheeres, D. [2003] Stability of relative equilibria in the full two-body problem. *Annals of the New York Academy of Science* This collection.
4. Wisdom, J. [1987] Rotational dynamics of irregularly shaped natural satellites. *Astron. J.* **94**:1350–1360.
5. Touma, J. and Wisdom, J. [1994] Evolution of the Earth-Moon system. *Astron. J.* **108**:1943–1961.

6. Beck, J. and Hall, C. [1998] Relative equilibria of a rigid satellite in a circular keplerian orbit. *J. Astronautical Sciences* **46**:215–247.
7. Wang, L. S., Krishnaprasad, P. S., and Maddocks, J. H. [1991] Hamiltonian dynamics of a rigid body in a central gravitational field. *Celest. Mech. Dyn. Astron.* **50**:349–386.
8. Wang, L. S., Krishnaprasad, P. S., and Maddocks, J. H. [1992] Steady rigid-body motions in a central gravity field. *J. Astronautical Sciences* **40**:449–478.
9. Wang, L. S., Lian, K.-Y., and Chen, P.-T. [1995] Steady motions of gyrostat satellites and their stability. *IEEE Trans. Automatic Control* **40**:1732–1743.
10. Rui, C., Kolmanovsky, I., and McClamroch, N. H. [2000] Nonlinear attitude and shape control of spacecraft with articulated appendages and reaction wheels. *IEEE Trans. Automatic Control* **45**:1455–1469.
11. Shen, J. and McClamroch, N. H. [2001] Translational and rotational spacecraft maneuvers via shape change actuators. In *Proceedings of 2001 American Control Conference*, 3961–3966.
12. Bloch, A. M., Crouch, P., Marsden, J. E., et al. [2002] The symmetric representation of the rigid body equations and their discretization. *Nonlinearity* **15**:1309–1341.
13. Dobrovolskis, A. and Burns, J. [1980] Life near the Roche limit: Behavior of ejecta from satellites close to planets. *Icarus* **42**:422–441.
14. B., C., P., F., and F., M. [1993] Planar orbits about a triaxial body: Applications to asteroidal satellites. *Icarus* **105**:370–384.
15. D.J., S. [1994] Dynamics About Uniformly Rotating Tri-Axial Ellipsoids. Applications to Asteroids. *Icarus* **110**:225–238.
16. J., S. D., J., O. S., S., H. R., et al. [1996] Orbits close to asteroid 4769 Castalia. *Icarus* **121**:67–87.
17. D.J., S., S.J., O., R.S., H., et al. [1998] Dynamics of orbits close to asteroid 4179 Toutatis. *Icarus* **132**:53–79.
18. Scheeres, D., Durda, D., and Geissler, P. [2003] The fate of asteroid ejecta. In *Asteroids III*. Univ. of Arizona Press, Tuscon. In press.
19. Scheeres, D., B.G., W., and J.K., M. [2000] Evaluation of the dynamic environment of an asteroid: Applications to 433 Eros. *J. of Guidance, Control and Dynamics* **23**:466–475.
20. Duboshin, G. [1958] The Differential Equations of Translational - Rotational Motion of Mutually Attracting Rigid Bodies. *Astronomicheskii Zhurnal* **35**:265.
21. Maciejewski, A. J. [1995] Reduction, relative equilibria and potential in the two rigid bodies problem. *Celest. Mech. Dyn. Astron.* **63**:1–28.
22. Scheeres, D. [2002] Stability in the full two-body problem. *Celest. Mech. Dyn. Astron.* **83**:155–169.
23. D.J., S., S.J., O., R.A., W., et al. [2000] Effects of gravitational interactions on asteroid spin states. *Icarus* **147**:106–118.
24. Scheeres, D. [2001] Changes in rotational angular momentum due to gravitational interactions between two finite bodies. *Celest. Mech. Dyn. Astron.* **81**:39–44.
25. Scheeres, D. [2002] Stability of Binary Asteroids. *Icarus* **159**:271–283.

26. Bloch, A. [2003] *Nonholonomic Mechanics and Control*. Applied Mathematical Sciences Series. Springer-Verlag.
27. Marsden, J. E. and Ratiu, T. S. [1999] *Introduction to Mechanics and Symmetry*, vol. 17 of *Texts in Applied Mathematics*. Springer-Verlag.
28. Marsden, J. [1992] *Lectures on Mechanics*, vol. 174 of *London Math. Soc. Lecture Note Ser.*. Cambridge University Press.
29. Bloch, A. M., Krishnaprasad, P. S., Marsden, J. E., et al. [1994] Dissipation induced instabilities. *Ann. Inst. H. Poincaré, Analyse Nonlinéaire* **11**:37–90.
30. Bloch, A. M., Krishnaprasad, P. S., Marsden, J. E., et al. [1996] The Euler–Poincaré equations and double bracket dissipation. *Comm. Math. Phys.* **175**:1–42.
31. Duncan, M., Quinn, T., and Tremaine, S. [1989] The long-term evolution of orbits in the solar system - A mapping approach. *Icarus* **82**:402–418.
32. Wisdom, J. and Holman, M. [1991] Symplectic maps for the n -body problem. *Astron. J.* **102**:1528–1538.
33. Lew, A., Marsden, J. E., Ortiz, M., et al. [2003] Variational time integration for mechanical systems. *Intern. J. Num. Meth. in Engin.* To appear.
34. Lew, A., Marsden, J. E., Ortiz, M., et al. [2003] Asynchronous variational integrators. *Arch. Rat. Mech. Anal.* To appear.
35. Jaffé, C., Ross, S. D., Lo, M. W., et al. [2002] Statistical Theory of Asteroid Escape Rates. *Phys. Rev. Lett.* **89**:011101–1.
36. Dellnitz, M., Junge, O., Koon, W.-S., et al. [2003] Transport in dynamical astronomy and multibody problems Preprint.
37. Néron de Surgy, O. and Laskar, J. [1997] On the long term evolution of the spin of the Earth. *Astronomy and Astrophysics* **318**:975–989.
38. Greenberg, R., Hoppa, G., Geissler, P., et al. [2002] The rotation of Europa. *Celest. Mech. Dyn. Astron.* **83**:35–47.
39. Bottke, W., Jedicke, R., Morbidelli, A., et al. [2000] Understanding the distribution of near-Earth asteroids. *Science* **288**:2190–2194.
40. Bottke, W. F., Vokrouhlický, D., Brož, M., et al. [2001] Dynamical Spreading of Asteroid Families by the Yarkovsky Effect. *Science* **294**:1693–1696.
41. Spitale, J. and Greenberg, R. [2001] Numerical evaluation of the general Yarkovsky effect: Effects on semimajor axis. *Icarus* **149**:222–234.
42. D., R. [2000] Radiative Spin-up and Spin-down of Small Asteroids. *Icarus* **148**:2–11.
43. Neishtadt, A., Scheeres, D., Sidorenko, V., et al. [2002] Evolution of comet nucleus rotation. *Icarus* **157**:205–218.
44. Yeomans, D. and Chodas, P. [1989] An asymmetric outgassing model for cometary nongravitational accelerations. *Astron. J.* **98**:1083–1093.
45. Margot, J. L., Nolan, M. C., Benner, L. A. M., et al. [2002] Binary Asteroids in the Near-Earth Object Population. *Science* **296**:1445–1448.

46. Veillet, C., Parker, J., Griffin, I., et al. [2002] The binary Kuiper-belt object 1998 WW31. *Nature* **416**:711.
47. Merline, W. J., Weidenschilling, S. J., Durda, D. D., et al. [2002] Asteroids Do Have Satellites. In *Asteroids III* (edited by W. Bottke, A. Cellino, P. Paolicchi, and R. P. Binzel), 289–312. Univ. of Arizona Press, Tucson.
48. Belton, M., Mueller, B., D’Amario, L., et al. [1996] The discovery and orbit of 1993 (243)1 Dactyl. *Icarus* **120**:185–199.
49. Benner, L., Ostro, S., Giorgini, J., et al. [2001] 1999 KW4. *IAU Circular* **7730**.
50. Harris, A. [2002] On the slow rotation of asteroids. *Icarus* **156**:184–190.
51. D.J., S. [1999] The Effect of C_{22} on Orbit Energy and Angular Momentum. *Celest. Mech. Dyn. Astron.* **73**:339–348.
52. Simo, J. C., Lewis, D. R., and Marsden, J. E. [1991] Stability of relative equilibria I: The reduced energy momentum method. *Arch. Rat. Mech. Anal.* **115**:15–59.
53. Marsden, J. E. and West, M. [2001] Discrete mechanics and variational integrators. *Acta Numerica* **10**:357–514.
54. Koon, W., Lo, M., Marsden, J., et al. [2000] Heteroclinic connections between periodic orbits and resonance transitions in celestial mechanics. *Chaos* **10**:427–469.
55. Cendra, H., Marsden, J. E., and Ratiu, T. S. [2001] *Lagrangian reduction by stages*, vol. 152 of *Memoirs*. American Mathematical Society, Providence, R.I.
56. Marsden, J. E., Ratiu, T. S., and Scheurle, J. [2000] Reduction theory and the Lagrange-Routh equations. *J. Math. Phys.* **41**:3379–3429.
57. Maciejewski, A. J. [1999] The two rigid bodies problem. Reduction and relative equilibria. In *Hamiltonian systems with three or more degrees of freedom (S’Agaró, 1995)*, 475–479.
58. Goździewski, K. and Maciejewski, A. J. [1999] Unrestricted planar problem of a symmetric body and a point mass. Triangular libration points and their stability. *Celest. Mech. and Dyn. Astron.* **75**:251–285.
59. Leonard, N. E. and Marsden, J. E. [1997] Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry. *Physica D* **105**:130–162.
60. Marsden, J., Misiolek, G., Perlmutter, M., et al. [1998] Symplectic reduction for semidirect products and central extensions. *Diff. Geom. and its Appl.* **9**:173–212.
61. Marsden, J. E., Montgomery, R., and Ratiu, T. S. [1990] *Reduction, symmetry and phases in mechanics*, vol. 436 of *Memoirs of the AMS, vol 436*. Amer. Math. Soc., Providence, RI.
62. Clerc, M. and Marsden, J. E. [2001] Dissipation-induced instabilities in an optical cavity laser: A mechanical analog near the 1:1 resonance. *Physical Rev. E* **64**:067603.
63. Lara, M. and Scheeres, D. [2003] Stability bounds for three-dimensional motion close to asteroids. *J. of the Astronautical Sciences* In press.
64. Alber, M., Luther, G., Marsden, J., et al. [1998] Geometric phases, reduction and Lie-Poisson structure for the resonant three-wave interaction. *Physica D* **123**:271–290.
65. Hernandez, A. and Marsden, J. E. [2002] Bifurcation of singular relative equilibria by the blowing up method Submitted for publication.

66. Stern, S., Bottke, W., and Levison, H. [2003] Regarding the putative eccentricity of Charon's orbit. *Astron. J.* **125**:902–905.
67. Sawai, S., Scheeres, D., and Broschart, S. [2002] Control of Hovering Spacecraft using Altimetry. *J. of Guidance, Control, and Dynamics* **25**:786–795.
68. Chang, D., Bloch, A. M., Leonard, N., et al. [2002] The equivalence of controlled Lagrangian and controlled Hamiltonian systems. *Control and the Calculus of Variations (special issue dedicated to J.L. Lions)* **8**:393–422.
69. Rom-Kedar, V. and Wiggins, S. [1990] Transport in two-dimensional maps. *Arch. Rat. Mech. Anal.* **109**:239–298.
70. Lekien, F. and Marsden, J. [2003] Separatrices in high-dimensional phase spaces: application to Van Der Waals dissociation In preparation.
71. Lekien, F. and Coulliette, C. [2003] MANGEN: Computation of hyperbolic trajectories, invariant manifolds and lobes of dynamical systems defined as 2D+1 data sets In preparation.
72. Lekien, F. [2003] *Time-Dependent Dynamical Systems and Geophysical Flows*. Ph.D. thesis, California Institute of Technology.
73. Wiggins, S. [1992] *Chaotic transport in dynamical systems*, vol. 2 of *Interdisciplinary Appl. Math.*. Springer, Berlin-Heidelberg-New York.
74. Rom-Kedar, V., Leonard, A., and Wiggins, S. [1990] An analytical study of transport, mixing and chaos in an unsteady vortical flow. *J. Fluid Mech.* **214**:347–394.
75. Rom-Kedar, V. and Wiggins, S. [1991] Transport in two-dimensional maps: Concepts, examples, and a comparison of the theory of Rom-Kedar and Wiggins with the Markov model of MacKay, Meiss, Ott, and Percival. *Physica D* **51**:248–266.
76. Coulliette, C. and Wiggins, S. [2001] Intergyre transport in a wind-driven, quasigeostrophic double gyre: an application of lobe dynamics. *Nonlinear Processes in Geophysics* **8**:69–94.
77. Lekien, F., Coulliette, C., and Marsden, J. [2003] Lagrangian structures in high-frequency radar data and optimal pollution timing. In *Proceedings of the 7th Experimental Chaos Conference*. Ameri. Inst. Phys. To appear.
78. Tiyapan, A. and Jaffé, C. [1995] Chaotic scattering: An invariant fractal tiling of phase space. *J. Chem. Phys.* **103**:5499–5511.
79. Jalnapurkar, S. M., Leok, M., Marsden, J. E., et al. [2003] Discrete Routh reduction. *Found. Comput. Math.* To appear.
80. Saha, P. and Tremaine, S. [1992] Symplectic integrators for solar system dynamics. *Astron. J.* **104**:1633–1640.
81. Marsden, J. E., Patrick, G. W., and Shkoller, S. [1998] Multisymplectic geometry, variational integrators and nonlinear PDEs. *Comm. Math. Phys.* 351–395.
82. Kane, C., Marsden, J., Ortiz, M., et al. [2000] Variational integrators and the Newmark algorithm for conservative and dissipative mechanical systems. *Int. J. Num. Meth. Eng.* **49**:1295–1325.
83. Rowley, C. W. and Marsden, J. E. [2002] Variational integrators for point vortices. *Proc. CDC* **40**:1521–1527.
84. Schlick, T. [2002] *Molecular modeling and simulation*, vol. 21 of *Interdisciplinary Applied Mathematics*. Springer-Verlag, New York.

85. Marsden, J. E., Pekarsky, S., and Shkoller, S. [2000] Symmetry reduction of discrete Lagrangian mechanics on Lie groups. *J. Geom. and Phys.* **36**:140–151.
86. Dellnitz, M. and Junge, O. [2002] Set oriented numerical methods for dynamical systems. In *Handbook of dynamical systems*, vol. 2, 221–264. North-Holland, Amsterdam.
87. Murray, C. and Dermott, S. [1999] *Solar System Dynamics*. Cambridge.
88. Cho, S., McClamroch, H., and Reyhanoglu, M. [2000] Dynamics of multi-body vehicles and their formulation as nonlinear control systems. In *Proceedings of 2000 American Control Conference*, 3908–3912.
89. Hagerty, P., Bloch, A., and Weinstein, M. [2003] Radiation induced instability in interconnected systems. In *Proc. of the 38th Conf. on Decisions and Control, IEEE*. To appear.
90. Washabaugh, P. and Scheeres, D. [2002] Energy and stress distributions in ellipsoids. *Icarus* **159**:314–321.
91. Asphaug, E. and Benz, W. [1996] Size, Density, and Structure of Comet Shoemaker-Levy 9 Inferred from the Physics of Tidal Breakup. *Icarus* **121**:225–248.
92. Richardson, D. C., Bottke, W. F., and Love, S. G. [1998] Tidal distortion and disruption of Earth-crossing asteroids. *Icarus* **134**:47–76.
93. Harris, A. [1994] Tumbling asteroids. *Icarus* **107**:209–211.
94. Bottke, W. F. and Melosh, H. J. [1996] The formation of binary asteroids and doublet craters. *Icarus* **124**:372–391.
95. Durda, D. D. [1996] The formation of asteroidal satellites in catastrophic collisions. *Icarus* **120**:212–219.
96. A., D., P., P., A., V., et al. [1997] The formation of binary asteroids as outcomes of catastrophic collisions. *Planet. Space. Sci.* **45**:757–770.
97. Bottke, W. F., Richardson, D. C., Michel, P., et al. [1999] 1620 Geographos and 433 Eros: Shaped by planetary tides? *Astron. J.* **117**:1921–1928.
98. Michel, P., Benz, W., Tanga, P., et al. [2001] Collisions and gravitational reaccumulation: Forming asteroid families and satellites. *Science* **294**:1696–1700.
99. Goldreich, P., Lithwick, Y., and Sari, R. [2002] Formation of Kuiper-belt binaries by dynamical friction and three-body encounters. *Nature* **240**:643–646.
100. Chandrasekhar, S. [1969] *Ellipsoidal Figures of Equilibrium*. Yale University Press.
101. Bloch, A. M., Chang, D., Leonard, N., et al. [2001] Controlled Lagrangians and the stabilization of mechanical systems II: Potential shaping. *IEEE Trans. Automatic Control* **46**:1556–1571.
102. Shen, J., McClamroch, N., and Bloch, A. [2003] Local equilibrium controllability of multibody systems controlled via shape change. *IEEE Trans. Automatic Control* To appear.
103. Meiss, J. [1992] Symplectic maps, variational principles, and transport. *Rev. Mod. Phys.* **64**:795–848.
104. Rom-Kedar, V. [1999] Transport in a class of n -d.o.f. systems. In *Hamiltonian systems with three or more degrees of freedom (S'Agaró, 1995)*, 538–543.

105. Beigie, D., Leonard, A., and Wiggins, S. [1991] A global study of enhanced stretching and diffusion in chaotic tangles. *Phys. Fluids A* **3**:1039–1049.
106. Malhotra, N. and Wiggins, S. [1998] Geometric structures, lobe dynamics, and Lagrangian transport in flows with aperiodic time dependence, with applications to Rossby wave flow. *J. of Nonlinear Science* **8**:401–456.
107. Poje, A. C. and Haller, G. [1999] Geometry of cross-stream mixing in a double-gyre ocean model. *Phys. Oceanogr.* **29**:1649–1665.
108. Haller, G. and Yuan, G. [2000] Lagrangian coherent structures and mixing in two-dimensional turbulence. *Physica D* **147**:352–370.
109. Haller, G. [2001] Distinguished material surfaces and coherent structures in 3D fluid flows. *Physica D* **149**:248–277.
110. Jaffé, C., Farrelly, D., and Uzer, T. [2000] Transition state theory without time-reversal symmetry: chaotic ionization of the hydrogen atom. *Phys. Rev. Lett.* **84**:610–613.
111. Ross, S., Koon, W., Lo, M., et al. [2003] Design of a Multi-Moon Orbiter. In *13th AAS/AIAA Space Flight Mechanics Meeting*. Ponce, Puerto Rico. AAS 03-143.