Chaos in the forced pendulum—an easy case of Lagrangian Coherent Structures

Philip DuToit and Jerrold E. Marsden
CDS, Caltech

CALTECH
Control & Dynamical Systems

March 15, 2007

Invariant Manifolds: Standard View

Invariant Manifolds: Standard View

- Start with the simple pendulum—a swing!

Invariant Manifolds: Standard View

- Start with the simple pendulum—a swing!

$$\ddot{x} + \sin x = 0$$

Invariant Manifolds: Standard View

- Start with the simple pendulum—a swing!

$$\ddot{x} + \sin x = 0$$

- Phase portrait—showing the invariant manifolds (separatrices attached to fixed points.

Invariant Manifolds: Standard View

- Start with the simple pendulum—a swing!

$$\ddot{x} + \sin x = 0$$

- Phase portrait—showing the invariant manifolds (separatrices attached to fixed points.



Homoclinic Chaos: Standard View

Homoclinic Chaos: Standard View

- Periodically perturb the simple pendulum with forcing

Homoclinic Chaos: Standard View

- Periodically perturb the simple pendulum with forcing

$$\ddot{x} + \sin x + \epsilon \dot{x} \sin t = 0$$

Homoclinic Chaos: Standard View

- Periodically perturb the simple pendulum with forcing

$$\ddot{x} + \sin x + \epsilon \dot{x} \sin t = 0$$

- Velocity field—hard to tell what is going on:

Homoclinic Chaos: Standard View

- Periodically perturb the simple pendulum with forcing

$$\ddot{x} + \sin x + \epsilon \dot{x} \sin t = 0$$

- Velocity field—hard to tell what is going on:



Standard way around this

Standard way around this

- Use of the Poincaré map (1880) to get a homoclinic tangle: excellent way to view for periodic systems.

Standard way around this

- Use of the Poincaré map (1880) to get a homoclinic tangle: excellent way to view for periodic systems.



Transient Chaos

Transient Chaos

- Poincaré's homoclinic tangle corresponds to transient chaos—dynamic events over intermediate time scales.

Transient Chaos

- Poincaré's homoclinic tangle corresponds to transient chaos—dynamic events over intermediate time scales.
- Infinite time notions like strange attractors, inertial manifolds, etc are not relevant in this context

Transient Chaos

- Poincaré's homoclinic tangle corresponds to transient chaos—dynamic events over intermediate time scales.
- Infinite time notions like strange attractors, inertial manifolds, etc are not relevant in this context
- First, a bit more about the tangle

Smale Horseshoe

Smale Horseshoe

- Smale abstracted what was going on in the tangle

Smale Horseshoe

- Smale abstracted what was going on in the tangle









Smale Horseshoe

- Smale abstracted what was going on in the tangle









- Proved lots of nice things—eg, an invariant Cantor set.

Smale Horseshoe in the Tangle

Smale Horseshoe in the Tangle



Smale Horseshoe in the Tangle



Smale Horseshoe in the Tangle







Smale Horseshoe in the Tangle









Smale Horseshoe in the Tangle



Smale Horseshoe in the Tangle



Lagrangian Coherent Structures

Lagrangian Coherent Structures

- Generalizes invariant manifolds to the case of time dependent dynamical systems

Lagrangian Coherent Structures

- Generalizes invariant manifolds to the case of time dependent dynamical systems
- For time varying systems, LCS move in time

Lagrangian Coherent Structures

- Generalizes invariant manifolds to the case of time dependent dynamical systems
- For time varying systems, LCS move in time



Look at lobes, mixing, dynamically