

CALIFORNIA INSTITUTE OF TECHNOLOGY  
Control and Dynamical Systems

**ACM/CDS 202**

R. Murray  
Spring 2013

**Problem Set #1**

Issued: 5 Apr 2013  
Due: 15 Apr 2013

Reading: Abraham, Marsden, and Ratiu, Sections 1.1–1.6

Problems:

1. (a) Consider  $[0, 2\pi)$  and  $\mathbb{S}^1 = \{x \in \mathbb{R}^2 \mid \|x\| = 1\}$  as subsets of  $\mathbb{R}$  and  $\mathbb{R}^2$ , respectively, and equip these sets with the relative topology. Define a map  $f : [0, 2\pi) \rightarrow \mathbb{S}^1$  as

$$f(x) = (\cos x, \sin x).$$

Is  $f$  a homeomorphism? Why or why not?

- (b) Let  $S$  be a set. Is the identity map from  $S$  with the discrete topology to  $S$  with the trivial topology continuous? a homeomorphism?
2. Show that a continuous bijection from a compact space to a Hausdorff space is always a homeomorphism. [*Hint*: Use the fact that a compact subspace of a Hausdorff space is closed.]
3. (a) Are the rational numbers a closed subset of  $\mathbb{R}$ ? Why or why not?  
(b) The set of rational numbers,  $\mathbb{Q}$ , is a subset of  $\mathbb{R}$  and hence inherits the usual metric on  $\mathbb{R}$  to become a metric space itself. Is it a complete metric space?
4. Let  $f : X \rightarrow \mathbb{R}$  be a continuous map and let  $X$  be compact. Show that  $f$  is bounded. That is, show that there exists  $M > 0$  such that  $|f(x)| \leq M$  for every  $x \in X$ .
5. [Abraham, Marsden, and Ratiu, Exercise 1.2-3]  
Show that every separable metric space is second countable.